

GAUGE TRANSFORMATIONS AND SYMMETRIES OF INTEGRABLE SYSTEMS

TAKESHI FUKUYAMA, KIYOSHI KAMIMURA, SAŠA KREŠIĆ-JURIĆ,
STJEPAN MELJANAC

ABSTRACT. We analyze several integrable systems in zero-curvature form within the framework of $SL(2, \mathbb{R})$ invariant gauge theory. In the Drinfeld-Sokolov gauge we derive a two-parameter family of nonlinear evolution equations which as special cases include the Kortweg-de Vries (KdV) and Harry Dym equations. We find residual gauge transformations which lead to infinitesimal symmetries of this family of equations. For KdV and Harry Dym equations we find an infinite hierarchy of such symmetry transformations, and we investigate their relation with local conservation laws, constants of the motion and the bi-Hamiltonian structure of the equations. Applying successive gauge transformations of Miura type we obtain a sequence of gauge equivalent integrable systems, among them the modified KdV and Calogero KdV equations.

1. INTRODUCTION

Integrable systems in $1 + 1$ dimensions have been discussed extensively so far, and various techniques to analyze their integrability have been developed using the Lax or zero-curvature representation, the inverse scattering method, the Hirota's bilinear method, the Painlevé property, etc. [1]. Integrable systems have also been investigated within the framework of factorization problems on loop groups and infinite dimensional Grassmannian spaces [2]. Gauge theories play a particular role in the investigation of integrable systems. In $1 + 1$ and higher dimensions there is a well-known conjecture by Ward [3] that any integrable systems are derived from (anti) self-dual Yang-Mills equations.

In this paper we examine several integrable systems from the point of view of $SL(2, \mathbb{R})$ invariant gauge theory. We consider infinitesimal gauge transformations acting on $sl(2, \mathbb{R})$ -valued fields which satisfy the zero-curvature equation. We analyze the transformations which leave

the given gauge fixing conditions invariant (i.e. residual transformations), and thus generate infinitesimal symmetries of evolution equations. Recall that if $u(x, t)$ is a solution of the evolution equation $u_t = K[u]$ where $K[u]$ depends on x , u , and a finite number of x -derivatives of u , then an infinitesimal symmetry is a transformation $u(x, t) \mapsto \tilde{u}(x, t) = u(x, t) + \tau y(x, t)$ such that the deformed function $\tilde{u}(x, t)$ is also a solution of the given equation to first order in τ . At least formally the infinitesimal transformations allow one to generate new solutions from old ones. Symmetries of evolution equations have also been studied by prolonged group actions on jet spaces, as discussed in Olver [4]. In this approach a local group of transformations acts on the space of independent and dependent variables, whereas gauge transformations considered here are globally defined groups acting on the space of dependent variables. For a particular class of evolution equations we investigate the relationship between residual gauge transformations, conservation laws and bi-Hamiltonian structure. We show that the conservation laws and the bi-Hamiltonian structure of the Korteweg-de Vries (KdV) and Harry Dym (HD) equations are directly related to an infinite hierarchy of residual transformations, and we thus provide a new interpretation of some previously known results from a unified perspective. We also investigate the Miura transformations from the gauge theoretical point of view. Since we do not incorporate the spectral parameter, i.e. we do not consider the loop algebra $sl(2, \mathbb{C}) \otimes \mathbb{C}[\lambda, \lambda^{-1}]$, the class of evolution equations obtained here is different from that considered in [5].

The outline of the paper is as follows. In Sec. 2 we consider the Drinfeld-Sokolov gauge on the Lie algebra $sl(2, \mathbb{R})$ and we derive a two-parameter family of nonlinear evolution equations which include as special cases the well-known KdV and HD equations. We find residual gauge transformations which lead to infinitesimal symmetries of this family of equations. A different derivation of KdV as a reduction of the self-dual Yang-Mills equations on \mathbb{R}^4 was given in [6] (see also [7]). In Sec. 3 we examine more closely the residual transformations for the KdV equation. We prove the existence of an infinite hierarchy of such transformations, and we show that each transformation generates

a local conservation law. We further show that the residual transformations are related to an infinite hierarchy of first integrals found in [8] and the Lenard recursion operator for the bi-Hamiltonian structure of KdV. Furthermore, by applying successively the Miura-type gauge transformations to the KdV equation we obtain a sequence of gauge equivalent integrable systems, among them the modified KdV (mKdV) and Calogero KdV (CKdV) equations. By using residual transformations we find nonlocal symmetries of the mKdV equation. Similar considerations are applied to the Harry Dym equation in Sec. 4. Finally, in Sec. 5 we note that by changing the gauge fixing condition and generalizing the gauge group to $SL(2, \mathbb{C})$ one obtains other classical integrable systems such as the sine-Gordon and nonlinear Schrödinger equation.

2. GENERAL FRAMEWORK

Let G be a matrix Lie group with Lie algebra \mathfrak{g} . Consider an overdetermined linear system of equations for a wave function ψ ,

$$(1) \quad \frac{\partial \psi}{\partial x_k} = A_k \psi, \quad 1 \leq k \leq m,$$

where $A_k(x) \in \mathfrak{g}$ is an $n \times n$ matrix valued gauge field and $\psi(x)$ is a vector valued function of $x \in \mathbb{R}^m$. The consistency condition $\partial_k \partial_l \psi = \partial_l \partial_k \psi$ requires that the gauge fields satisfy the zero-curvature condition [9]

$$(2) \quad \frac{\partial A_k}{\partial x_l} - \frac{\partial A_l}{\partial x_k} + [A_k, A_l] = 0, \quad 1 \leq k, l \leq m.$$

Equation (2) represents a hierarchy of nonlinear partial differential equations (PDEs) for components of the gauge fields. Equations which can be represented in this form are often said to be integrable, although the proof of integrability in each particular case is a nontrivial problem. The zero-curvature representation is used as a basis for application of geometric and algebraic techniques in the study of nonlinear PDEs, and it is also used in classification of integrable systems by Lie algebraic methods (see e.g. [10] and [11]). The Lie algebras considered here are finite dimensional, although a number of integrable systems can be written in zero-curvature form on infinite dimensional loop algebras. In this approach the equation at hand is integrated by solving

a Riemann-Hilbert type factorization problem on the underlying loop group (see e.g. [12]).

The system of equations (1)-(2) is invariant under the gauge transformation

$$(3) \quad \psi \mapsto g\psi,$$

$$(4) \quad A_k \mapsto \Gamma_g A_k \equiv g A_k g^{-1} + \frac{\partial g}{\partial x_k} g^{-1},$$

where $g \in G$. In general, this transformation changes the particular form of the equations represented by (2). Two systems of equations are said to be gauge equivalent if their zero-curvature representations are related by a gauge transformation. For examples of gauge equivalent systems see [10]. Gauge transformations can be used to map solutions of one system into the other [13]. In this paper we shall consider solutions of the system (1)-(2) which are gauge equivalent. Such solutions have the pure gauge form

$$(5) \quad \psi = g\psi_0, \quad A_k = \frac{\partial g}{\partial x_k} g^{-1},$$

for some constant vector $\psi_0 \in \mathbb{C}^n$ and $g \in G$. We note that, as pointed out by Wu and Yang [14], in some cases there exist gauge fields which give rise to the same field strength but are not related by any gauge transformations.

The zero-curvature condition (2) gives rise to integrable systems of a fairly general nature. However, one is usually interested in systems involving a smaller number of functions. This is achieved by imposing gauge fixing conditions which lead to a reduced set of differential equations for independent field components. In the following we discuss integrable systems in $1 + 1$ space-time dimensions for $sl(2, \mathbb{R})$ -valued fields. By parameterizing the gauge fields explicitly the linear problem becomes

$$(6) \quad \frac{\partial \psi}{\partial x} = A_1 \psi, \quad \frac{\partial \psi}{\partial t} = A_2 \psi.$$

where

$$(7) \quad A_1 = \begin{pmatrix} R & S \\ -T & -R \end{pmatrix}, \quad A_2 = \begin{pmatrix} p & u \\ q & -p \end{pmatrix}.$$

Here $\psi = (\psi_1 \ \psi_2)^t$, and the minus sign for T is introduced for convenience. The zero-curvature condition (2) yields

$$(8) \quad R_t - p_x = -qS - uT,$$

$$(9) \quad S_t - u_x = 2pS - 2uR,$$

$$(10) \quad T_t + q_x = -2qR - 2pT,$$

where the subscripts denote partial derivatives. There are three gauge freedoms for A_1 and A_2 since the six field components are related by three differential equations. First, we fix two of them by requiring

$$(11) \quad R = 0 \quad \text{and} \quad S = 1.$$

This is called the Drinfeld-Sokolov (DS) gauge [5]. In this gauge the fields are given by

$$(12) \quad A_1^{(DS)} = \begin{pmatrix} 0 & 1 \\ -T & 0 \end{pmatrix}, \quad A_2^{(DS)} = \begin{pmatrix} -\frac{1}{2}u_x & u \\ -\frac{1}{2}u_{xx} - uT & \frac{1}{2}u_x \end{pmatrix},$$

and the zero-curvature equation for $A_1^{(DS)}$ and $A_2^{(DS)}$ is equivalent to

$$(13) \quad T_t = 2u_x T + uT_x + \frac{1}{2}u_{xxx}.$$

We are interested in gauge transformations which preserve the gauge fixing condition (11), and thus leave equation (13) invariant. The existence of such residual transformations leads to symmetries of evolution equations and to an infinite hierarchy of conservation laws.

Define the map $K: sl(2, \mathbb{R}) \rightarrow \mathbb{R}^2$ by $K\left[\begin{pmatrix} a & b \\ c & -a \end{pmatrix}\right] = (a, b - 1)$, and consider the subvariety $\mathcal{S}_K = \{A \in sl(2, \mathbb{R}) \mid K(A) = 0\}$ which consists precisely of the matrices in the DS gauge. We note that K is of maximal rank since the Jacobian of K is

$$(14) \quad J_K = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}.$$

Fix $L \in sl(2, \mathbb{R})$ and consider the action of the one-parameter gauge group $G = \{\exp(\tau L) \mid \tau \in \mathbb{R}\}$ on $sl(2, \mathbb{R})$ by the gauge transformation (4). Since K is of maximal rank, G is a symmetry group of the set $\mathcal{S}_K \subset sl(2, A)$ if and only if each component of K satisfies (see e.g. [4])

$$(15) \quad \left. \frac{d}{d\tau} \right|_{\tau=0} K_i(\Gamma_{\exp(\tau L)} A) = 0 \quad \text{for all} \quad A \in \mathcal{S}_K, \quad i = 1, 2.$$

If L satisfies condition (15), then the gauge transformation $A_1^{(DS)} \mapsto \Gamma_g A_1^{(DS)}$ preserves the DS gauge and hence equation (13). In order to find L let $\pi: sl(2, \mathbb{R}) \rightarrow \mathbb{R}^2$ denote the projection $\pi \left[\begin{pmatrix} a & b \\ c & -a \end{pmatrix} \right] = (a, b)$. Then

$$(16) \quad \left. \frac{d}{d\tau} \right|_{\tau=0} K(\Gamma_{\exp(\tau L)} A) = \pi \left(\left. \frac{d}{d\tau} \right|_{\tau=0} \Gamma_{\exp(\tau L)} A \right),$$

where the expression in the parentheses defines the infinitesimal gauge transformation by L given by

$$(17) \quad \left. \frac{d}{d\tau} \right|_{\tau=0} \Gamma_{\exp(\tau L)} A = [L, A] + \frac{\partial L}{\partial x}.$$

Hence, in view of the condition (15) the DS gauge is preserved if and only if for every matrix A in the DS gauge, $A \in \mathcal{S}_K$, we have

$$(18) \quad \pi \left([L, A] + \frac{\partial L}{\partial x} \right) = 0.$$

Setting $A = A_1^{(DS)}$ and evaluating condition (18) we see that L can be parameterized by an arbitrary function $y(x, t)$ as

$$(19) \quad L = \begin{pmatrix} -\frac{1}{2}y_x & y \\ -\frac{1}{2}y_{xx} - yT & \frac{1}{2}y_x \end{pmatrix}.$$

The gauge transformation of $A_1^{(DS)}$ is now given by

$$(20) \quad \begin{aligned} \Gamma_{\exp(\tau L)} A_1^{(DS)} &= A_1^{(DS)} + \tau \left([L, A_1^{(DS)}] + \frac{\partial L}{\partial x} \right) + o(\tau^2) \\ &= \begin{pmatrix} 0 & 1 \\ -T & 0 \end{pmatrix} + \tau \begin{pmatrix} 0 & 0 \\ -\frac{1}{2}y_{xxx} - 2y_x T - yT_x & 0 \end{pmatrix} + o(\tau^2), \end{aligned}$$

hence the infinitesimal transformation of T is

$$(21) \quad \tilde{T} = T + \tau \left(\frac{1}{2}y_{xxx} + 2y_x T + yT_x \right).$$

Similarly, the gauge transformation of $A_2^{(DS)}$ yields

$$\begin{aligned}
 \Gamma_{\exp(\tau L)} A_2(DS) &= A_2^{(DS)} + \tau \left([L, A_2^{(DS)}] + \frac{\partial L}{\partial t} \right) + o(\tau^2) \\
 (22) \quad &= \begin{pmatrix} -\frac{1}{2}u_x & u \\ -\frac{1}{2}u_{xx} - uT & \frac{1}{2}u_x \end{pmatrix} \\
 &+ \tau \begin{pmatrix} \frac{1}{2}u_{xx}y + \frac{1}{2}uy_{xx} - \frac{1}{2}y_{xt} & y_t - y_xu + yu_x \\ \frac{1}{2}u_xy_{xx} - \frac{1}{2}u_{xx}y_x + (yu_x - y_xu)T - \frac{1}{2}y_{xxt} - (yT)_t & -\frac{1}{2}u_{xx}y - \frac{1}{2}uy_{xx} + \frac{1}{2}y_{xt} \end{pmatrix} \\
 &+ o(\tau^2).
 \end{aligned}$$

Thus, the infinitesimal transformation of u is given by

$$(23) \quad \tilde{u} = u + \tau(y_t - y_xu + yu_x).$$

For convenience we denote the infinitesimal deformations by $\delta T = \frac{1}{2}y_{xxx} + 2y_xT + yT_x$ and $\delta u = y_t - y_xu + yu_x$. Since the gauge transformations of $A_1^{(DS)}$ and $A_2^{(DS)}$ satisfy the zero-curvature equation (which can be also checked directly to first order in τ using equations (20) and (22)), the functions $\tilde{T} = T + \tau\delta T$ and $\tilde{u} = u + \tau\delta u$ satisfy equation (13) to first order in τ . Hence, the maps $T \mapsto \tilde{T}$ and $u \mapsto \tilde{u}$ are infinitesimal symmetries of (13).

The remaining gauge freedom is fixed by imposing a relation between T and u . We set

$$(24) \quad T = \frac{u^\alpha}{s}, \quad \alpha \in \mathbb{Z} \setminus \{0\}, \quad s > 0.$$

The zero-curvature condition (13) then becomes

$$(25) \quad u_t = \frac{\alpha + 2}{\alpha} u u_x + \frac{s}{2\alpha} u^{1-\alpha} u_{xxx}.$$

The motivation for considering the gauge (24) is that (25) is a two-parameter family of evolution equations which, as special cases, includes some well-known integrable systems: the KdV equation for $\alpha = 1$, $s = 2$, and Harry Dym equation for $\alpha = -2$, $s = 1$.

The infinitesimal symmetry of (25) is easily found by requiring that the function y is chosen so that \tilde{T} and \tilde{u} satisfy condition (24) to first order in τ . This holds if and only if

$$(26) \quad \delta T = \frac{\alpha}{s} u^{\alpha-1} \delta u.$$

By substituting (21) and (23) into (26) we find that y is constrained by the equation

$$(27) \quad y_t = \frac{\alpha + 2}{\alpha} u y_x + \frac{s}{2\alpha} u^{1-\alpha} y_{xxx}.$$

In this case the infinitesimal transformation for u is given by

$$(28) \quad \tilde{u} = u + \tau \left(\frac{2}{\alpha} u y_x + u_x y + \frac{s}{2\alpha} u^{1-\alpha} y_{xxx} \right).$$

Any smooth solution of equation (27) generates a residual gauge transformation of (25).

We noted earlier that equation (25) is integrable for $\alpha = 1, -2$. Then equation (27) has infinitely many solutions which generate a hierarchy of conservation laws for (25), as discussed later. Furthermore, a common property shared by all equations (25) is the existence of travelling wave solutions. We conjecture that equation (25) is integrable for all $\alpha \in \mathbb{Z} \setminus \{0\}$, which is certainly true if (27) has an infinite number of solutions for all such α . For a discussion of integrability of (25) using the Painlevé property see [15]. It is also noted that for $\alpha = 1$ the matrices $A_1^{(DS)}$ and $A_2^{(DS)}$ can be extended to matrix polynomials in the loop algebra $sl(2, \mathbb{C}) \otimes \mathbb{C}[\lambda, \lambda^{-1}]$ appearing in the Riemann-Hilbert factorization for the KdV equation (see [12]). It would be a very interesting problem to determine if this construction carries over for all α and if (25) can be integrated by solving a Riemann-Hilbert factorization problem.

As stated earlier, equations (25) admit travelling wave solutions $u(x + vt)$. They are given explicitly for $\alpha = \pm 1, \pm 2$, and in implicit form for other values of α . The solutions are assumed to satisfy the boundary conditions $u, u', u'' \rightarrow 0$ as $|x + vt| \rightarrow \infty$. For $\alpha = 2k - 1$, $k \in \mathbb{N}$, the solution is given implicitly by

$$(29) \quad \sqrt{1 - bu} F\left(\frac{1}{2}, k, \frac{3}{2}; 1 - bu\right) = \frac{a}{2b^{k-1}}(x + vt + x_0)$$

where

$$(30) \quad a = \sqrt{\frac{4v}{s(\alpha + 1)}}, \quad b = \frac{\alpha}{sv},$$

and F is the hypergeometric function [16]

$$(31) \quad F(\alpha, \beta, \gamma; z) = \sum_{k=0}^{\infty} \frac{(\alpha)_k (\beta)_k}{(\gamma)_k} \frac{z^k}{k!}.$$

Similarly, for $\alpha = 2k$, $k \in \mathbb{N}$, we find

$$(32) \quad \sqrt{1-bu}F\left(\frac{1}{2}, \frac{1}{2} - k, \frac{3}{2}; 1-bu\right) = \frac{a}{2b^{k-\frac{1}{2}}}(x+vt+x_0).$$

For $\alpha = 1$ (and $s = 2$) equation (29) yields the well-known one-soliton solution for KdV,

$$(33) \quad u(x, t) = 2v \operatorname{sech}^2\left(\sqrt{\frac{v}{4}}(x+vt+x_0)\right),$$

while for $\alpha = 2$ it gives a rational solution

$$(34) \quad u(x, t) = \frac{4}{a^2(x+vt+x_0)^2 + 4b}.$$

The travelling wave solutions for negative integers α are given by similar expressions. For $\alpha = -(2k+1)$, $k \in \mathbb{N}$, we have

$$(35) \quad \sqrt{1-bu}F\left(\frac{1}{2}, -k, \frac{3}{2}; 1-bu\right) = \frac{ab^{k+1}}{2}(x+vt+x_0),$$

and for $\alpha = -2k$, $k \in \mathbb{N} \setminus \{1\}$, we find

$$(36) \quad \sqrt{1-bu}F\left(\frac{1}{2}, \frac{1}{2} - k, \frac{3}{2}; 1-bu\right) = \frac{ab^{k+\frac{1}{2}}}{2}(x+vt+x_0).$$

For $\alpha = -2$ (and $s = 1$) we obtain a rational solution of the Harry Dym equation,

$$(37) \quad u(x, t) = (9v)^{\frac{1}{3}}(x+vt+x_0)^{\frac{2}{3}}.$$

When $\alpha = -1$ the solution is given as a series $u(x, t) = \sum_{n=0}^{\infty} c_n(x+vt)^n$. The coefficients c_n are found recursively from

$$(38) \quad \sum_{k=0}^{\infty} \alpha_k \beta_{n-k} + \frac{2v}{s}(n-2)c_{n-2} - \frac{2}{s^2}\gamma_{n-2} = 0, \quad n \geq 3,$$

where α_n , β_n , γ_n are defined by

$$(39) \quad \alpha_n = n(n-1)(n-2)c_n, \quad \beta_n = \sum_{k=0}^n c_{n-k}c_k, \quad \gamma_n = \sum_{k=0}^n kc_{n-k}c_k.$$

The recurrence relation places no restrictions on c_0 , c_1 and c_2 . For $n \geq 3$ the coefficient c_n is uniquely determined from c_0, c_1, \dots, c_{n-1} , hence the solution u is nontrivial provided c_0 , c_1 and c_2 are all nonzero. The existence of travelling wave solutions is a consequence of the spatio-temporal translational symmetry of equation (25). Although the travelling wave solutions are not directly related to integrability of (25), they may prove helpful in investigating the system's properties.

3. GAUGE TRANSFORMATIONS AND THE KdV SEQUENCES

In this section we consider more closely the residual gauge transformations for the KdV equation and their relation with the Lenard recursion operator for the bi-Hamiltonian structure of KdV. We also investigate gauge transformations between the KdV and related systems, as well as their residual transformations. If $T = u/2$ ($\alpha = 1$, $s = 2$) the gauge fields in (12) become

$$(40) \quad A_1^{(K)} = \begin{pmatrix} 0 & 1 \\ -\frac{1}{2}u & 0 \end{pmatrix}, \quad A_2^{(K)} = \begin{pmatrix} -\frac{1}{2}u_x & u \\ -\frac{1}{2}u_{xx} - \frac{1}{2}u^2 & \frac{1}{2}u_x \end{pmatrix},$$

hence (25) yields

$$(41) \quad u_t = 3uu_x + u_{xxx}.$$

This is one of the equivalent forms of the KdV equation. Equation (27) gives the residual gauge transformation for the KdV equation,

$$(42) \quad y_t = 3uy_x + y_{xxx}.$$

This is the linear equation associated with the KdV equation which Gardner et. al. obtained from the inverse scattering method [17]. If y is a solution of (42), then $\tilde{u} = u + \tau\delta u$ is an infinitesimal transformation of the KdV equation, where according to (28)

$$(43) \quad \delta u(y) = 2uy_x + u_x y + y_{xxx}.$$

Equation (42) has an infinite number of solutions for the parameter function y which are related to a hierarchy of conservation laws for the KdV equation. Let $y^{(1)}$ be a solution of (42) and define the function

$$(44) \quad G^{(1)} \equiv \delta u(y^{(1)}) = 2u y_x^{(1)} + u_x y^{(1)} + y_{xxx}^{(1)}.$$

Then $G^{(1)}$ satisfies the evolution equation

$$(45) \quad G_t^{(1)} = G_{xxx}^{(1)} + 3u_x G^{(1)} + 3u G_x^{(1)}.$$

This can be seen by replacing the time derivatives in $G_t^{(1)}$ by (41) and (42), and expressing $G_t^{(1)}$ only in terms of the x -derivatives of u and $y^{(1)}$. We note that equation (45) can be written as a local conservation law

$$(46) \quad \frac{\partial G^{(1)}}{\partial t} = \frac{\partial F^{(1)}}{\partial x}$$

with density $G^{(1)}$ and flux $F^{(1)} = G_{xx}^{(1)} + 3u G^{(1)}$. Now consider the function $y^{(2)}$ defined by $y_x^{(2)} = G^{(1)}$. The conservation law (46) implies that time evolution of $y^{(2)}$ is given by

$$(47) \quad y_t^{(2)} = \int G_t^{(1)} dx = 3u y_x^{(2)} + y_{xxx}^{(2)},$$

which is of the same form as (42). Hence, $y^{(2)}$ also generates a residual gauge transformation for KdV. By iterating the above procedure we obtain an infinite hierarchy of residual transformations $y^{(n)}$ defined recursively by

$$(48) \quad y_x^{(n+1)} = G^{(n)}, \quad G^{(n)} = \delta u(y^{(n)}), \quad n \in \mathbb{N},$$

together with local conservation laws

$$(49) \quad \frac{\partial G^{(n)}}{\partial t} = \frac{\partial F^{(n)}}{\partial x}, \quad F^{(n)} = G_{xx}^{(n)} + 3u G^{(n)}, \quad n \in \mathbb{N}.$$

To illustrate the point, we start with the simplest solution $y^{(1)} = 1$. Then the first four residual transformations yield

$$(50) \quad y^{(1)} = 1,$$

$$(51) \quad y^{(2)} = u,$$

$$(52) \quad y^{(3)} = \frac{3}{2}u^2 + u_{xx},$$

$$(53) \quad y^{(4)} = \frac{15}{6}u^3 + \frac{5}{2}u_x^2 + 5u u_{xx} + u_{xxxx}.$$

We remark that the functions $y^{(n)}$ are related to the isospectral problem for the Schrödinger equation, as shown in [18]. One can prove that if the eigenvalue λ in the Schrödinger equation $\Psi_{xx} + \frac{1}{6}u\Psi = \lambda\Psi$ is t -independent, then u must satisfy the evolution equation $u_t = \delta u(y^{(n)})$ for some $n \in \mathbb{N}$. In this way starting from the isospectral problem for the Schrödinger operator one can associate to KdV a hierarchy of integrable equations. A different approach to derive integrable hierarchies using Sato's pseudo differential operator was given in [19].

Next we show that the functions $y^{(n)}$ are related to an infinite hierarchy of conserved quantities for the KdV equation as follows. Let $u_t = K[u]$ be an evolution equation for $u(x, t)$ and suppose that the functional

$$(54) \quad \mathcal{H}_n[u] = \int_{-\infty}^{\infty} H_n[u] dx, \quad n \geq 1,$$

is a conserved quantity under the flow $u_t = K[u]$. In [8] Wadati showed that the variational derivative of \mathcal{H}_n satisfies the integral equation

$$(55) \quad \int_{-\infty}^{\infty} \left(h(x) \frac{\partial}{\partial t} \left(\frac{\delta \mathcal{H}_n}{\delta u} \right) + dK(u, h) \frac{\delta \mathcal{H}_n}{\delta u} \right) dx = 0$$

for every function $h(x)$ with compact support, where $dK(u, h)$ is the differential $dK(u, h) = \frac{d}{d\tau} \Big|_{\tau=0} K(u + \tau h)$. If we take $K[u] = 3u u_x + u_{xxx}$, then (55) implies that $\delta \mathcal{H}_n / \delta u$ satisfies equation (42). Thus, to each conserved quantity \mathcal{H}_n of the KdV equation we can associate a residual transformation by

$$(56) \quad y^{(n)} = \frac{\delta \mathcal{H}_n}{\delta u}.$$

It is worth noting that there is a close connection between the functions $G^{(n)}$, $y^{(n)}$ and the Lenard recursion operator for the bi-Hamiltonian structure of KdV [4]. Indeed, in view of (43) we have $G^{(n)} = E y^{(n)}$ where E is the differential operator $E = D^3 + 2uD + u_x$ and D denotes the derivative with respect to x . Then (48) implies

$$(57) \quad G^{(n)} = R G^{(n-1)},$$

where $R = ED^{-1}$ is precisely the Lenard operator for the KdV equation. We also note that

$$(58) \quad y^{(n)} = R^* y^{(n-1)}$$

where $R^* = D^{-1}E$ is the adjoint of R . This implies that the Hamiltonians \mathcal{H}_n defined by (56) are related by the Lenard recursion formula

$$(59) \quad D \frac{\delta \mathcal{H}_n}{\delta u} = E \frac{\delta \mathcal{H}_{n-1}}{\delta u}.$$

Solving equation (56) for \mathcal{H}_2 and \mathcal{H}_3 we obtain

$$(60) \quad \mathcal{H}_2[u] = \int_{-\infty}^{\infty} \frac{1}{2} u^2 dx, \quad \mathcal{H}_3[u] = \int_{-\infty}^{\infty} \left(-\frac{1}{2} u_x^2 + \frac{1}{2} u^3 \right) dx.$$

The Hamiltonians \mathcal{H}_2 and \mathcal{H}_3 give rise to the well-known bi-Hamiltonian structure for KdV since

$$(61) \quad u_t = D \frac{\delta \mathcal{H}_3}{\delta u} = E \frac{\delta \mathcal{H}_2}{\delta u}.$$

All Hamiltonians \mathcal{H}_n are conserved under the KdV flow, and they are all in involution with each other with respect to the Gardner-Poisson brackets for the Hamiltonian operators D and E [20]. Hence, equation (48) can be viewed as another interpretation of the bi-Hamiltonian

property of the KdV equation. Applications of gauge transformations to Hamiltonian systems from a different point of view can be found in [21].

Next we consider the Miura transformation from the gauge theoretical point of view. Miura-type transformations are known to provide links between different integrable systems, as discussed in [22]. Recall that with the choice $T = u/2$ there is no gauge freedom left for the matrices $A_1^{(K)}$ and $A_2^{(K)}$ in equation (40), and the zero-curvature condition is equivalent with the KdV equation (41). Consider the gauge transformation of $A_1^{(K)}$ and $A_2^{(K)}$ defined by the matrix

$$(62) \quad g = \begin{pmatrix} 1 & 0 \\ -v & 1 \end{pmatrix}$$

for some $v = v(x, t)$. The gauge fields transform according to

$$(63) \quad A_1^{(M)} = g A_1^{(K)} g^{-1} + g_x g^{-1} = \begin{pmatrix} v & 1 \\ -\frac{1}{2}u - v^2 - v_x & -v \end{pmatrix},$$

$$(64) \quad A_2^{(M)} = g A_2^{(K)} g^{-1} + g_t g^{-1} = \begin{pmatrix} uv - \frac{1}{2}u_x & u \\ -\frac{1}{2}u^2 - \frac{1}{2}u_{xx} - v^2u + v u_x - v_t & -uv + \frac{1}{2}u_x \end{pmatrix}.$$

Now suppose that v and u are related by the Miura transformation

$$(65) \quad v^2 + v_x = -\frac{1}{2}u.$$

Then $A_1^{(M)}$ and $A_2^{(M)}$ become

$$(66) \quad A_1^{(M)} = \begin{pmatrix} v & 1 \\ 0 & -v \end{pmatrix}, \quad A_2^{(M)} = \begin{pmatrix} -2v^3 + v_{xx} & -2(v^2 + v_x) \\ -v_t - 6v^2v_x + v_{xxx} & 2v^3 - v_{xx} \end{pmatrix},$$

and the zero-curvature condition is equivalent with

$$(67) \quad \left(\frac{\partial}{\partial x} + 2v \right) (-v_t - 6v^2v_x + v_{xxx}) = 0.$$

Since the zero-curvature equation is invariant under gauge transformations, we recover the well-known result which states that if v satisfies the modified KdV (mKdV) equation

$$(68) \quad v_t = v_{xxx} - 6v^2v_x,$$

then $u = -2(v^2 + v_x)$ solves the KdV equation. Note that the reverse implication does not necessarily hold.

By using the same arguments as in Sec. 2 one can find the residual transformations for the mKdV equation. Define $K: sl(2, \mathbb{R}) \rightarrow \mathbb{R}^2$ by $K\left(\begin{smallmatrix} a & b \\ c & -a \end{smallmatrix}\right) = (b-1, c)$, and consider the subvariety

$$(69) \quad \mathcal{S}_K = \{A \in sl(2, \mathbb{R}) \mid K(A) = 0\}$$

which consists of matrices of the form $A_1^{(M)}$. Define the gauge group $G = \{\exp(\tau M) \mid \tau \in \mathbb{R}\}$ for some $M \in sl(2, \mathbb{R})$. Then G is a symmetry group of the set \mathcal{S}_K if and only if the infinitesimal gauge transformation by M satisfies

$$(70) \quad \pi \left([M, A] + \frac{\partial M}{\partial x} \right) = 0 \quad \text{for all } A \in \mathcal{S}_K,$$

where the projection $\pi: sl(2, \mathbb{R}) \rightarrow \mathbb{R}^2$ is given by $\pi\left(\begin{smallmatrix} a & b \\ c & -a \end{smallmatrix}\right) = (b, c)$. Evaluation of the condition (70) with $A = A_1^{(M)}$ shows that M can be parameterized by a single function $z(x, t)$ as

$$(71) \quad M = \begin{pmatrix} z v - \frac{1}{2} z_x & z \\ c & -z v + \frac{1}{2} z_x \end{pmatrix} \quad \text{where } c = \exp \left(-2 \int v dx \right),$$

Now, the gauge transformation of $A_1^{(M)}$ yields

$$(72) \quad \Gamma_{\exp(\tau M)} A_1^{(M)} = A_1^{(M)} + \tau \left([M, A_1^{(M)}] + \frac{\partial M}{\partial x} \right) + o(\tau^2)$$

$$(73) \quad = \begin{pmatrix} v & 1 \\ 0 & -v \end{pmatrix} + \tau \begin{pmatrix} -\frac{1}{2} z_{xx} + (zv)_x - c & 0 \\ 0 & \frac{1}{2} z_{xx} - (zv)_x + c \end{pmatrix} + o(\tau^2),$$

hence

$$(74) \quad \tilde{v} = v + \tau \left(-\frac{1}{2} z_{xx} + (zv)_x - c \right)$$

is an infinitesimal transformation of the mKdV equation. This transformation is nonlocal because c is defined in terms of an indefinite integral. Among the first to consider such transformations and to point out some applications were Vinogradov and Krasilshchik [23]. Recently, nonlocal symmetries for some well-known integrable systems were found in [24].

We are left with the problem of determining the evolution equation for the function z . Let $u = -2(v^2 + v_x)$ denote the $(1, 2)$ element of the matrix $A_2^{(M)}$ in equation (66). The gauge transformation of $A_2^{(M)}$ yields the infinitesimal transformation of u ,

$$(75) \quad \tilde{u} = u + \tau (z_t - u z_x + u_x z).$$

Since \tilde{u} and \tilde{v} are also related by $\tilde{u} = -2(\tilde{v}^2 + \tilde{v}_x)$ to first order in τ , this implies that z satisfies the equation

$$(76) \quad z_t = z_{xxx} - 6(v^2 + v_x) z_x.$$

We note that this is just the residual symmetry of the KdV equation for the function $u = -2(v^2 + v_x)$. Therefore, if z is a solution of equation (76), then $v \mapsto \tilde{v} = v + \tau \delta v$ is an infinitesimal symmetry of the mKdV equation where

$$(77) \quad \delta v = -\frac{1}{2} z_{xx} + (z v)_x - \exp \left(-2 \int v dx \right).$$

The gauge transformation with g of the form (62) can be repeatedly applied to obtain other integrable systems. Consider the transformations of $A_1^{(M)}$ and $A_2^{(M)}$ defined by the matrix

$$(78) \quad h = \begin{pmatrix} 1 & 0 \\ -w & 1 \end{pmatrix}.$$

We have

$$(79) \quad A_1^{(C)} = h A_1^{(M)} h^{-1} + h_x h^{-1} = \begin{pmatrix} v + w & 1 \\ -2v w - w^2 - w_x & -(v + w) \end{pmatrix}$$

$$(80) \quad A_2^{(C)} = h A_2^{(M)} h^{-1} + h_t h^{-1}$$

$$(81) \quad = \begin{pmatrix} -2v^3 - 2(v^2 + v_x) w + v_{xx} & -2(v^2 + v_x) \\ 2(2v^3 - v_{xx}) w + 2(v^2 + v_x) w^2 - w_t & 2v^3 + 2(v^2 + v_x) w - v_{xx} \end{pmatrix}.$$

If we choose w so that the $(2, 1)$ element of $A_1^{(C)}$ vanishes, i.e.

$$(82) \quad w_x + w^2 + 2vw = 0,$$

then the zero-curvature condition for $A_1^{(C)}$ and $A_2^{(C)}$ yields the Calogero KdV (CKdV) equation [25]

$$(83) \quad w_t = w_{xxx} - \frac{1}{2} (3w_x^2 w^{-1} + w^3)_x.$$

Thus, by successively applying the gauge transformations of the form (62) we obtain a sequence of integrable equations called the KdV sequence. The successive Bäcklund transformations from the KdV to CKdV equation were discussed in the bilinear formalism in [26]. The same sequence of integrable equations was obtained by Pavlov [27] by considering solutions of the overdetermined system

$$(84) \quad \psi_{xx} = \left(\lambda - \frac{1}{2}u\right)\psi, \quad \psi_t = (4\lambda + u)\psi_x - \frac{1}{2}u_x\psi,$$

whose consistency condition yields the KdV equation. He showed that by expanding the wave function into a power series in the spectral parameter λ ,

$$(85) \quad \psi = \exp \left(\int \left(\sum_{n=0}^{\infty} r^{(n)} \lambda^n \right) dx \right),$$

the coefficients $r^{(n)}(x, t)$ satisfy the same sequence of equations derived above. Thus, the successive transformations with no spectral parameter in our theory correspond to the expansion of the wave function into power series in the spectral parameter in conventional theories.

4. THE HARRY DYM EQUATION

The Harry Dym (HD) equation [28]

$$(86) \quad u_t = -\frac{1}{4}u^3 u_{xxx}$$

is obtained from (25) by setting $\alpha = -2$ and $s = 1$. It was discovered by Harry Dym when trying to transfer some results about the isospectral theory to the string equation [29]. The HD equation is a completely integrable Hamiltonian system which is solvable by the inverse scattering transform [30], [31], and it possesses the bi-Hamiltonian structure and infinitely many conservation laws [32].

In this case the fields in (12) become

$$(87) \quad A_1^{(HD)} = \begin{pmatrix} 0 & 1 \\ -\frac{1}{u^2} & 0 \end{pmatrix}, \quad A_2^{(HD)} = \begin{pmatrix} -\frac{1}{2}u_x & u \\ -\frac{1}{2}u_{xx} - \frac{1}{u} & \frac{1}{2}u_x \end{pmatrix}.$$

In view of equations (27) and (28) the infinitesimal symmetry transformation for the HD equation is given by

$$(88) \quad \tilde{u} = u + \tau(-u y_x + u_x y - \frac{1}{4}u^3 y_{xxx}),$$

where y is constrained by the equation

$$(89) \quad y_t = -\frac{1}{4}u^3 y_{xxx}.$$

We show that equation (89) has an infinite number of solutions for the parameter function y . Suppose $y^{(1)}$ is a solution of (89) and define $y^{(2)}$ by

$$(90) \quad \left(\frac{y^{(2)}}{u}\right)_x = -u y_{xxx}^{(1)}.$$

We claim that $y^{(2)}$ is also a solution of equation (89). Denote for the moment $y_{xxx}^{(1)} = v$ so that $y^{(2)} = -u \int uv dx$. Then a simple computation shows that

$$(91) \quad y_t^{(2)} = \frac{1}{4}u^3 u_{xxx} \int uv dx + \frac{1}{4}u \left(\int u_{xxx} u^3 v dx + \int u(u^3 v)_{xxx} dx \right)$$

where we used the assumption that $y^{(1)}$ satisfies (89), and we replaced the time derivative of u by equation (86). After a few partial integrations of the last term we obtain

$$(92) \quad \int u(u^3 v)_{xxx} dx = u(u^3 v)_{xx} - u_x(u^3 v)_x + u_{xx}u^3 v - \int u_{xxx}u^3 v dx,$$

which finally yields

$$(93) \quad y_t^{(2)} = \frac{1}{4}u^3 \left(u_{xxx} \int uv dx + 3u_x^2 v + 5uu_x v_x + 4uu_{xx} v + u^2 v_{xx} \right).$$

It is straightforward to verify that the expression in the parentheses equals $-y_{xxx}^{(2)}$, hence $y^{(2)}$ satisfies equation (89). Therefore, (90) provides a recursion formula for $y^{(n)}$ which generates an infinite hierarchy of residual gauge transformations for the HD equation. Starting with $y^{(1)} = 1$ the first four functions are given by

$$(94) \quad y^{(1)} = 1,$$

$$(95) \quad y^{(2)} = u,$$

$$(96) \quad y^{(3)} = \frac{1}{2}u u_x^2 - u^2 u_{xx},$$

$$(97) \quad y^{(4)} = \frac{3}{8}u u_x^4 - \frac{3}{2}u^2 u_x^2 u_{xx} + \frac{3}{2}u^3 u_{xx}^2 + 2u^3 u_x u_{xxx} + u^4 u_{xxxx}.$$

As in the case of the KdV equation the conserved quantities of the HD equation are related to the residual gauge transformations $y^{(n)}$. Let

$$(98) \quad \mathcal{G}_n[u] = \int_{-\infty}^{\infty} G_n[u] dx, \quad n \in \mathbb{N},$$

be a conserved quantity of the HD equation (86). Substituting $K[u] = -\frac{1}{4}u^3u_{xxx}$ into equation (55) one can show that $\delta\mathcal{G}_n/\delta u$ satisfies

$$(99) \quad \left(\frac{\delta\mathcal{G}_n}{\delta u} \right)_t = \frac{3}{4}u^2u_{xxx} \frac{\delta\mathcal{G}_n}{\delta u} - \frac{1}{4} \left(u^3 \frac{\delta\mathcal{G}_n}{\delta u} \right)_{xxx}.$$

We emphasize that, unlike in the KdV case, the above equation is not the same as the residual transformation (89) for $y^{(n)}$. However, if we set

$$(100) \quad y^{(n)} = u^3 \frac{\delta\mathcal{G}_n}{\delta u},$$

then it is easily seen that $y^{(n)}$ satisfies (89) if and only if $\delta\mathcal{G}_n/\delta u$ satisfies (99). Therefore, the conserved quantities and residual gauge transformations of the HD equation are related by equation (100). The conserved quantities corresponding to the first four residual transformations $y^{(n)}$ found earlier are given by

$$(101) \quad \mathcal{G}_1[u] = \int_{-\infty}^{\infty} \left(-\frac{1}{2}u^{-2} \right) dx,$$

$$(102) \quad \mathcal{G}_2[u] = \int_{-\infty}^{\infty} (-u^{-1}) dx,$$

$$(103) \quad \mathcal{G}_3[u] = \int_{-\infty}^{\infty} \frac{1}{2}u^{-1}u_x^2 dx,$$

$$(104) \quad \mathcal{G}_4[u] = \int_{-\infty}^{\infty} \left(\frac{1}{8}u^{-1}u_x^4 + \frac{1}{2}uu_{xx}^2 \right) dx.$$

5. THE SINE-GORDON AND NONLINEAR SCHRÖDINGER EQUATIONS

So far we have discussed integrable systems on the Lie algebra $sl(2, \mathbb{R})$ which arise by fixing the DS gauge (11). In this section we consider briefly a few generalizations obtained by changing the gauge fixing condition and generalizing the gauge group. More precisely, we derive the sine-Gordon (SG) and focusing nonlinear Schrödinger (NLS) equations. The NLS equation was derived by symmetry reduction from the self-dual Yang-Mills equations in [6] (see also [7]). We remark that the residual transformations approach presented in the previous sections

can be extended to study infinitesimal symmetries of the SG and NLS equations as well.

Consider the gauge fields (7) with $R = \alpha$, $\alpha \in \mathbb{R} \setminus \{0\}$, and $S = T = -\frac{1}{2}w_x$ for some $w = w(x, t)$. In order to fix the remaining gauge freedom suppose $q = u$. Then the zero-curvature equations (8)-(10) yield

$$(105) \quad p_x = -q w_x,$$

$$(106) \quad \frac{1}{2}w_{xt} + q_x = p w_x + 2\alpha q,$$

$$(107) \quad -\frac{1}{2}w_{xt} + q_x = p w_x - 2\alpha q.$$

Adding and subtracting equations (106) and (107) we obtain

$$(108) \quad q_x = p w_x,$$

$$(109) \quad w_{xt} = 4\alpha q.$$

We note that equations (105) and (108) admit solutions $p = \cos(w)$ and $q = \sin(w)$. With this choice of p and q the gauge fields become

$$(110) \quad A_1^{(SG)} = \begin{pmatrix} \alpha & -\frac{1}{2}w_x \\ \frac{1}{2}w_x & -\alpha \end{pmatrix}, \quad A_2^{(SG)} = \begin{pmatrix} \cos(w) & \sin(w) \\ \sin(w) & -\cos(w) \end{pmatrix}.$$

The zero-curvature condition is then equivalent with (109), which is the sine-Gordon equation

$$(111) \quad w_{xt} = 4\alpha \sin(w).$$

Similarly, we derive the focusing NLS equation by considering the gauge group $SL(2, \mathbb{C})$ and introducing the gauge fixing conditions $R = 0$, $S = \bar{T} = w$ and $q = -\bar{u}$. In this case the zero-curvature equations (8)-(10) yield

$$(112) \quad -q w + \bar{q} \bar{w} = -p_x,$$

$$(113) \quad w_t + \bar{q}_x = 2p w,$$

$$(114) \quad \bar{w}_t + q_x = -2p \bar{w}.$$

Equation (112) implies that p_x is purely imaginary, hence we may impose further reduction of the gauge freedom by requiring $p = -i|w|^2$. Then equations (113) and (114) are complex conjugate of each other.

Furthermore, if we set $q = -i\overline{w}_x$, then relation (112) holds identically. In this case the gauge fields are

$$(115) \quad A_1^{(NLS)} = \begin{pmatrix} 0 & w \\ -\overline{w} & 0 \end{pmatrix}, \quad A_2^{(NLS)} = \begin{pmatrix} -i|w|^2 & -iw_x \\ -i\overline{w}_x & i|w|^2 \end{pmatrix},$$

and (114) yields the focusing NLS equation

$$(116) \quad iw_t - w_{xx} - 2|w|^2 w = 0.$$

The sine-Gordon and the focusing NLS equation can also be derived within the framework of Birkhoff factorization of loop groups. For more details see e.g. [13] and [33].

6. CONCLUSION

In this paper we have discussed integrable systems within the framework of gauge theory. The gauge theoretical structure of the Zakharov-Shabat formalism has been fully discussed. Starting with the DS gauge, we have fixed the residual gauge freedom by the relation (24) and obtained a two-parameter family of nonlinear equations (25). It was shown explicitly that for $\alpha = 1$ (KdV) and $\alpha = -2$ (HD) there exist infinite hierarchies of residual gauge transformations. The transformations can be calculated recursively and they lead to an infinite number of conserved quantities. We have also shown that all equations (25) admit travelling wave solutions which are given implicitly in terms of the hypergeometric function, except for $\alpha = \pm 1, \pm 2$ when the solutions are given in explicit form. However, it is an open problem to determine if equation (25) is integrable for all $\alpha \in \mathbb{Z} \setminus \{0\}$. The Miura transformations which lead to the KdV sequence (KdV \rightarrow mKdV \rightarrow CKdV, etc.) have been formulated in terms of gauge transformations acting on the gauge fields defined on the Lie algebra $sl(2, \mathbb{R})$. It was shown that other types of 1+1 integrable systems can also be formulated within the same framework by changing the initial gauge fixing condition and generalizing the gauge group.

REFERENCES

REFERENCES

- [1] Zakharov V E ed 1991 *What is Integrability?* (Berlin: Springer-Verlag)
- [2] Segal G and Wilson G 1985 *Publ. Math. I.H.E.S.* **61** 5–65

- [3] Ward R S 1985 *Phil. Trans. R. Soc. London, Ser. A* **315** 451–457
- [4] Olver P 1993 *Applications of Lie Groups to Differential Equations* 2nd ed (New York: Springer-Verlag)
- [5] Drinfeld V and Sokolov V 1984 *J. Sov. Math.* **30** 1975–2036
- [6] Mason L J and Sparling G A J 1989 *Phys. Lett. A* **137** 29–33
- [7] Ablowitz M J and Clarkson P A 1991 *Solitons, Nonlinear Evolution Equations and Inverse Scattering* (Cambridge: Cambridge University Press)
- [8] Wadati M 1978 *Stud. Appl. Math.* **59** 153–186
- [9] Zakharov V E and Shabat A B 1972 *JETP* **34** 62–69
- [10] Fadeev L D and Takhtajan T A 1987 *Hamiltonian Methods in the Theory of Solitons* (New York: Springer-Verlag)
- [11] Babelon O, Bernard D and Talon M 2003 *Introduction to Classical Integrable Systems* (Cambridge: Cambridge University Press)
- [12] Dorfmeister J 1993 *Proc. Symp. Pure Math.* **54** 121–139
- [13] Krešić-Jurić S 2006 *J. Math. Phys.* **47** 063501
- [14] Wu T T and Yang C N 1975 *Phys. Lett. D* **12** (12) 3843–3844
- [15] Fukuyama T, Kamimura K and Toda K nlin.SI/0108043, unpublished
- [16] Lebedev N N 1972 *Special Functions and Their Applications* (New York: Dover)
- [17] Gardner C S, Green J M, Kruskal M D and Miura R M 1974 *Comm. Pure Appl. Math.* **27** 97–133
- [18] Das A 1989 *Integrable Models* (Singapore: World Scientific)
- [19] Kundu A, Strampp W and Oevel W 1995 *J. Math. Phys.* **36** (6) 2972–2984
- [20] Lax P D 1976 *SIAM Review* **18** 351–375
- [21] Liu X, Chen Q and Zhang B 2003 *Appl. Math. Mech.* **24** (7) 863–870
- [22] Pelinovsky D, Springael J, Lambert F and Loris I 1997 *J. Phys. A: Math. Gen.* **30** 8705–8717
- [23] Vinogradov A M and Krasilshchik I S 1980 *Dokl. Akad. Nauk. SSSR* **253** (6) 1289–1293
- [24] Schiff J *Preprint* arxiv.org, solv-int/9606004
- [25] Calogero F and Degasperis A 1981 *J. Math. Phys.* **22** 23–31
- [26] Nakamura A and Hirota R 1980 *J. Phys. Soc. Japan* **48** 1755–1762
- [27] Pavlov M V 1998 *Phys. Lett. A* **243** 295–300
- [28] M.D. Kruskal M D 1975 *Lecture Notes in Physics* vol 38 (Berlin: Springer-Verlag)
- [29] Hereman W, Banerjee P P and Chatterjee M R 1989 *J. Phys. A: Math. Gen.* **22** 241–255
- [30] Wadati M, Ichikawa Y H and Shimizu T 1980 *Prog. Theor. Phys.* **64** 1959–1967
- [31] Wadati M, Konno K and Ichikawa Y H 1979 *J. Phys. Soc. Japan* **47** 1698–1700
- [32] Brunelli J C and da Costa G 2002 *J. Math. Phys.* **43** 6116–6128
- [33] Dorfmeister J and Wu H 1993 *J. Reine Angew. Math.* **440** 43–76

(T. Fukuyama) DEPARTMENT OF PHYSICS, RITSUMEIKAN UNIVERSITY, SHIGA,
KUSATSU 525-8577, JAPAN

(K. Kamimura) DEPARTMENT OF PHYSICS, TOHO UNIVERSITY, FUNABASHI
247-8510, JAPAN

(S. Krešić-Jurić) FACULTY OF NATURAL AND MATHEMATICAL SCIENCES, UNI-
VERSITY OF SPLIT, TESLINA 12, 21000 SPLIT, CROATIA

E-mail address: `skresic@fesb.hr`

(S. Meljanac) RUDJER BOŠKOVIĆ INSTITUTE, BIJENIČKA CESTA B.B., 10000
ZAGREB, CROATIA